



UNIVERSITI PUTRA MALAYSIA

**EFFECT OF RICE QUALITY, FORMULATION AND
STORAGE ON THE QUALITY OF CANNED RICE PORRIDGE**

MA YONG QIN

FSMB 2000 10

**EFFECT OF RICE QUALITY, FORMULATION AND
STORAGE ON THE QUALITY OF CANNED RICE PORRIDGE**

MA YONG QIN

**MASTER OF SCIENCE
UNIVERSITI PUTRA MALAYSIA**

2000



**EFFECT OF RICE QUALITY, FORMULATION AND
STORAGE ON THE QUALITY OF CANNED RICE PORRIDGE**

By

MA YONG QIN

**Thesis Submitted in Fulfilment of the Requirements for the
Degree of Master of Science in the Faculty of
Food Science and Biotechnology
Universiti Putra Malaysia**

May 2000



Dedicated to

*My beloved parents, husband, daughter,
sisters and brother*

*for their love,
patience
and understanding*

Abstract of the thesis presented to the Senate of Universiti Putra Malaysia in
fulfilment of the requirements for the degree of Master of Science

**EFFECT OF RICE QUALITY, FORMULATION AND
STORAGE ON THE QUALITY OF CANNED RICE PORRIDGE**

By

MA YONG QIN

May 2000

Chairman : Sharifah Kharidah Syed Muhammad, Ph.D.

Faculty : Food Science and Biotechnology

Long-grain rice was mixed with broken rice or glutinous rice at a ratio of 0:100, 25:75, 50:50, 75:25 and 100:0, respectively. Each mixture was processed into canned plain rice porridge. The physical and sensory properties of the products were evaluated. An optimum ratio was selected for the preparation of savoury rice porridge (chicken, beef, fish and bean flavours). The freshly prepared savoury rice porridges were then evaluated to determine their physicochemical, microbiological and sensory qualities. A storage study was also carried out to determine the quality changes of canned savoury rice porridges stored at 27°C and 4°C for 12 weeks.

The results obtained showed that plain rice porridge was best prepared using 100% long-grain rice. However, plain rice porridge prepared using a mixture of broken rice and long-grain rice at a ratio of 50:50 was found to be as acceptable as that containing 100% long-grain rice. Incorporation of glutinous rice reduced phase

separation but in excess of 25 percent led to a significant decrease in acceptability. Physical characteristics of the plain rice porridge were relatively unchanged with the substitution of long-grain rice with broken rice. When long-grain rice was substituted with increasing amounts of glutinous rice, the pH value and whiteness of the plain rice porridge tended to decrease but its %Brix and viscosity increased gradually. For the savoury rice porridges, the results of the ranking test revealed that the fish rice porridge was the most preferred followed by the beef, chicken and bean rice porridges in terms of colour, texture, flavour and taste. Results of the nutrient composition analyses showed that the beef rice porridge provided the highest protein, fat, and caloric contents and the most amounts of calcium and iron. Results of the storage study indicated that the temperature and time of storage did not affect dry matter, ash, protein, fat, caloric, calcium and iron content of the porridges. However, low storage temperature caused the porridges to have a lower %Brix and increase in whiteness. Porridges stored at room temperature were more stable in appearance, colour, flavour and taste than those stored in the refrigerator.

A composite of broken rice and long-grain rice at a ratio of 50:50 is recommended for the production of plain rice porridge. A can of the four types of savoury rice porridges designed can provide 1/4-1/5 of the recommended daily allowance (RDA) for energy and nutrients for 7-12 years old children. All samples stored at room temperature or in the refrigerator complied with the standard for food hygiene and safety up to 12 weeks of storage. However, canned rice porridge stored at room temperature gave better sensory characteristics.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Master Sains

**KESAN KUALITI BERAS, FORMULASI DAN PENYIMPANAN
KEATAS KUALITI BUBUR NASI DALAM TIN**

Oleh

MA YONG QIN

Mei 2000

Pengerusi : Dr. Sharifah Kharidah Syed Muhammad, Ph.D.

Fakulti : Sains Makanan dan Bioteknologi

Beras panjang dicampurkan dengan beras hancur atau beras pulut masing-masing pada nisbah 0:100, 25:75, 50:50, 75:25 dan 100:0. Setiap campuran telah dijadikan bubur nasi tawar dalam tin. Seterusnya, ciri-ciri fizikal dan deria produk-produk tersebut dinilai. Satu nisbah optima telah dipilih untuk penyediaan bubur nasi berperisa (perisa ayam, daging lembu, ikan dan kacang). Bubur nasi berperisa di nilai sejurus selepas disediakan untuk menentukan ciri-ciri fiziko-kimia, mikrobiologi dan derianya. Satu kajian penyimpanan juga telah dijalankan untuk menentukan perubahan kualiti bubur nasi berperisa dalam tin yang disimpan pada suhu 27°C dan 4°C selama 12 minggu.

Keputusan yang diperolehi menunjukkan bahawa bubur nasi tawar yang terbaik boleh disediakan menggunakan 100% beras panjang. Namun demikian bubur nasi tawar yang disediakan dengan campuran beras hancur dan beras panjang pada nisbah 50:50 dapat diterima setanding dengan bubur nasi tawar yang mengandungi

100% beras panjang. Penambahan beras pulut mengurangkan pemisahan fasa tetapi jika melebihi 25% mengakibatkan penerimaan bubur nasi tawar berkurangan secara bererti. Ciri fizikal produk didapati tidak berubah apabila beras panjang digantikan dengan beras hancur. Apabila beras panjang digantikan dengan kuantiti beras pulut yang semakin meningkat, nilai pH dan keputihan produk didapati berkurangan manakala %Brix dan kepekatan meningkat secara beransur-ansur. Bagi bubur nasi berperisa, keputusan ujian mengikut susunan keutamaan menunjukkan bahawa bubur nasi ikan paling disukai diikuti dengan bubur nasi daging lembu, bubur nasi ayam dan bubur nasi kekacang dari segi warna, tekstur, perisa dan rasa. Melalui analisis komposisi nutrien didapati yang bubur nasi daging lembu mempunyai kandungan protein, lemak, kalori, kalsium dan zat besi yang tertinggi. Keputusan kajian penyimpanan pula menunjukkan bahawa suhu dan masa penyimpanan tidak mempengaruhi kandungan bahan kering, abu, protein, lemak, kalori, kalsium dan besi dalam bubur nasi. Walau bagaimanapun, suhu penyimpanan yang rendah mengakibatkan bubur nasi mengalami pengurangan nilai %Brix dan penambahan kesan keputihan. Bubur nasi yang telah disimpan pada suhu bilik didapati lebih stabil dari segi rupa luaran, warna, perisa dan citarasa jika dibandingkan dengan bubur nasi yang telah disimpan dalam peti sejuk.

Campuran beras hancur dan beras panjang pada nisbah 50:50 disarankan untuk penghasilan bubur nasi tawar. Setiap tin daripada empat jenis bubur nasi berperisa yang dicipta akan memberikan 1/4-1/5 tenaga dan nutrien-nutrien yang disyorkan bagi kanak-kanak berusia di antara 7-12 tahun. Semua sampel yang

disimpan pada suhu bilik atau dalam peti sejuk memenuhi piawaian makanan bersih dan selamat sehingga jangka penyimpanan 12 minggu. Walau bagaimanapun, bubur nasi dalam tin yang disimpan pada suhu bilik telah memberikan ciri-ciri deria yang lebih baik.

ACKNOWLEDGEMENTS

I would like to express my most sincere appreciation and very deepest gratitude to Dr. Sharifah Kharidah Syed Muhammad for her kind understanding and willingness to be the chairman of my supervisory committee at a crucial part of this study. Her very generous gift of time and guidance, invaluable advice, constructive suggestions and comments had helped tremendously in the preparation of this thesis. I am also very grateful for the patience, care and encouragement that she had so generously offered during this study and in reviewing the thesis.

My heartfelt appreciation and sincere thanks go to the members of my supervisory committee, Prof. Madya Dr. Salmah Yusof and En. Dzulkifly Mat Hashim, who had selflessly devoted their time and reviewed the thesis in great detail time after time. I would also like to thank them for their very generous help, care, encouragement and warm friendship, which had made my study in UPM a very pleasant and memorable experience.

My deepest appreciation and special thanks go to my former supervisor, Dr. Asbi Bin Ali for his brilliant ideas, technical guidance and advice during this study.

My deepest appreciation and gratitude is accorded to all my friends, technicians, laboratory assistants and staff members of the Faculty of Food Science

and Biotechnology for their very generous help, care, encouragement and warm friendship.

My special thanks to my parents who are most dear to me for their heavenly affection and great sacrifice for my study and the other members of my family for their understanding, support and love.

I specially acknowledge my boundless gratitude to my husband for his constant support and encouragement, which enabled me toward completing the study.


I certify that an Examination Committee met on 16th May 2000 to conduct the final examination of Ma Yong Qin on her Master of Science thesis entitled "Effect of Rice Quality, Formulation and Storage on the Quality of Canned Rice Porridge" in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulations 1981. The committee recommends that the candidate be awarded the relevant degree. Members of the Examination Committee are as follows:

Jinap Selamat, Ph.D.
Professor,
Department of Food Science
Faculty of Food Science and Biotechnology
Universiti Putra Malaysia
(Chairman)

Sharifah Kharidah Syed Muhammad, Ph.D.
Associate Professor,
Department of Food Science
Faculty of Food Science and Biotechnology
Universiti Putra Malaysia
(Member)

Salmah bt. Yusof, Ph.D.
Associate Professor,
Department of Food Technology
Faculty of Food Science and Biotechnology
Universiti Putra Malaysia
(Member)

Dzulkifly Mat Hashim,
Lecturer
Department of Food Technology
Faculty of Food Science and Biotechnology
Universiti Putra Malaysia
(Member)



.....
MOHD. GHAZALI MOHAYIDIN, Ph.D.
Professor/Deputy Dean of Graduate School,
Universiti Putra Malaysia

Date: 29 MAY 2000

This thesis submitted to the Senate of Universiti Putra Malaysia and was accepted as fulfilment of the requirements for the degree of Master of Science.



KAMIS AWANG, Ph.D.
Associate Professor
Dean of Graduate School,
Universiti Putra Malaysia

Date: 13 JUL 2000

DECLARATION

I hereby declare that the thesis is based on my original work except for quotations and citations which have been duly acknowledged. I also declare that it has not been previously or concurrently submitted for any other degree at UPM or other institutions.



Ma Yong Qin

Date: 19/5/2001

TABLE OF CONTENTS

	Page
DEDICATION	2
ABSTRACT	3
ABSTRAK	5
ACKNOWLEDGEMENTS	8
APPROVAL SHEETS	10
DECLARATION	12
LIST OF TABLES	16
LIST OF FIGURES	18
LIST OF PLATES	19
CHAPTER	
I GENERAL INTRODUCTION	20
II LITERATURE REVIEW	24
Importance of Rice as Food	24
Trends in Consumption of Rice in Some Rice-eating Countries- Taiwan, Japan, Malaysia and Thailand	26
Rice-based Products in Some Rice-eating Countries-Taiwan, Japan, Malaysia and Thailand	28
Rice-based Products in Taiwan, Republic of China	28
Traditional and New Rice-based Products in Japan	30
Rice-base Products in Malaysia and Thailand	33
Definition of Rice Porridge	36
Classification of Rice Porridge	37
Quick-cooking Rice Porridge	37
Ready-to-eat Rice Porridge	42
Formulation of Rice Porridge	43
Processing of Rice Porridge	46
Processing of Quick-cooking Rice Porridge	46
Processing of Ready-to-eat Rice Porridge	48
Quality Characteristics of Ready-to-eat Rice Porridge	52
Sensory Quality	52
Physicochemical and Microbiological Characteristics	54
III EFFECTS OF VARIOUS COMPOSITE RICE PROPORTIONS BASED ON LONG GRAIN, BROKEN AND GLUTINOUS RICE ON PHYSICAL AND SENSORY CHARACTERISTICS OF CANNED PLAIN RICE PORRIDGE	56

Introduction	56
Materials and Methods	58
Materials	58
Experimental Design	59
Preparation of Canned Plain Rice Porridge	61
Analyses of Raw Rice	61
Determination of Physical Characteristics of Canned Plain Rice Porridge	64
Sensory Evaluation	66
Statistical Analysis	67
Results and Discussion	67
Quality Characteristics of Raw Rice	67
Pasting Behaviour of Raw Rice Flour	69
Effects of Composite Rice Proportions Based on Broken/Long Grain Rice and Glutinous/Long Grain Rice on pH, %Brix, Phase Separation and Viscosity of Canned Plain Rice Porridge	73
Canning Stability of Long-grain Rice, Broken Rice and Glutinous Rice	80
Colour of Raw Rice and Canned Plain Rice Porridge	84
Sensory Evaluation	88
Conclusion	90
 IV FORMULATION OF CANNED SAVOURY RICE PORRIDGE FOR PRIMARY SCHOOL CHILDREN	 91
Introduction	91
Materials and Methods	93
Materials	93
Sample Preparation	93
Physicochemical Analyses	98
Sensory Analysis	98
Statistical Analysis	99
Results and Discussion	99
Proximate Composition of Canned Savoury Rice Porridges	99
Mineral Content of Canned Savoury Rice Porridges.....	102
Physical Characteristics of Canned Savoury Rice Porridge.....	105
Sensory Evaluation	106
Conclusion	109

V	EFFECTS OF STORAGE CONDITIONS ON THE QUALITY OF CANNED SAVOURY RICE PORRIDGE	111
	Introduction	111
	Materials and Methods	112
	Materials	112
	Experimental Design	112
	Physicochemical Analyses	113
	Microbial Analysis	114
	Sensory Analysis	114
	Statistical Analysis	114
	Results and Discussion	115
	Physicochemical Changes in Rice Porridge During Storage	115
	Microbial Analysis	120
	Sensory Characteristics during Storage	122
	Conclusion	126
VI	GENERAL CONCLUSION AND RECOMMENDATIONS	128
	Conclusions	128
	Recommendations	129
	BIBLIOGRAPHY	131
	APPENDICES	
	A Analytical Methodology	138
	B Sensory Evaluation Sheets	148
	C Additional Table	152
	VITA	153

LIST OF TABLES

Table		Page
1	Commodity, Production and Price of Japanese Traditional Rice-based Foods (1986)	31
2	Commodity, Production and Price of New-processed Rice Foods in Japan (1986)	32
3	Common Rice-Based Products in Peninsular Malaysia	34
4	Traditional Rice-Based Products in Peninsular Malaysia	35
5	Processing and Characteristics of Flaked Rice Porridge	39
6	Types of Quick-cooking Rice Porridge	41
7	Type, Formulation, Processing and Packing Material of Some Ready-to-eat Rice Porridges.....	45
8	Quality Characteristics of Ready-to-eat Rice Porridge	53
9	Composition of Composite Rice Evaluated	59
10	Quality Characteristics of Raw Rice	68
11	Pasting Behaviour of Composite Rice Flour Based on Long-grain, Broken and Glutinous Rice	72
12	Correlation of Physicochemical Characteristics of Rice with Rice Porridge Quality	76
13	Analysis of Particle Size of Canned Plain Rice Porridge.....	83
14	Sensory Attributes of Canned Plain Rice Porridge	89
15	Formulation of Canned Chicken Rice Porridge (CP)	94
16	Formulation of Canned Beef Rice Porridge (BFP)	95
17	Formulation of Canned Fish Rice Porridge (FP)	95

18	Formulation of Canned Bean Rice Porridge (BNP)	96
19	Composition of Savoury Rice Porridges	100
20	Mineral Content of Savoury Rice Porridge	103
21	Physical Properties of Various Savoury Rice Porridges	106
22	Sensory Attributes of Various Savoury Rice Porridges	107
23	Rank Sum Analysis for Four Formulations	109
24	Dry Matter, Ash Content, pH Value and %Brix of Rice Porridges During Storage	116
25	Hunterlab L, a and b Values for Rice Porridges During Storage	119
26	Protein, Fat, Caloric, Calcium and Iron Contents of Rice Porridges Stored at 0 and 12 Weeks	121
27	Total Plate Count and Coliform of Rice Porridges During Storage	122
28	Sensory Scores for Appearance, Colour, Flavour and Taste of Rice Porridges During Storage Obtaining Using Quantitative Description Analysis	123
29	Sensory Scores for Appearance, Colour, Flavour and Taste of Rice Porridges During Storage Obtaining Using Hedonic Scale	124
30	Ranking of Four Formulations by 10 Panelists	152

LIST OF FIGURES

Figure		Page
1	Ready-to-eat Rice Porridge Canning Process	49
2	Preparation of Canned Plain Rice Porridge	60
3	Viscosity Profiles of Composite Rice Flour	70
4	Viscosity Profiles of Composite Rice Flour (Continued)	71
5	Effects of Composite Rice Proportions Based on Broken/Long Rice (B-L) and Glutinous/Long Rice(G-L) on pH values of Canned Plain Rice Porridge	74
6	Effects of Composite Rice Proportions Based on Broken/Long Rice (B-L) and Glutinous/Long Rice (G-L) on %Brix of Canned Plain Rice Porridge	77
7	Effects of Composite Rice Proportions Based on Broken/Long Rice (B-L) and Glutinous/Long Rice (G-L) on Viscosity of Canned Plain Rice Porridge	79
8	Effects of Composite Rice Proportions Based on Broken/Long Rice (B-L) and Glutinous/Long Rice (G-L) on Solids Loss of Canned Plain Rice Porridge	81
9	Effects of Composite Rice Proportions Based on Broken/Long Rice (B-L) and Glutinous/Long Rice (G-L) on Hunterlab L Value of Canned Plain Rice Porridge	85
10	Effects of Composite Rice Proportions Based on Broken/Long Rice (B-L) and Glutinous/Long Rice (G-L) on Hunterlab a Value of Canned Plain Rice Porridge	86
11	Effects of Composite Rice Proportions Based on Broken/Long Rice (B-L) and Glutinous/Long Rice (G-L) on Hunterlab b Value of Canned Plain Rice Porridge	87
12	Preparation of Canned Savoury Rice Porridge	97
13	Determination of Total Protein by Semi-micro Kjeldahl Method	141

LIST OF PLATES

Plate		Page
1	Rice Porridge Sold in Malaysian Market.....	37
2	Varieties of Quick-cooking Rice Porridge	40
3	Canned Sweet Rice Porridges Made in China	43

CHAPTER I

GENERAL INTRODUCTION

Rice porridge is a very popular food in the Asian region, particularly in China, Japan and Korea (Lu, 1987). It is usually home-made, and also available to consumers in the restaurants in freshly cooked form. Commercial rice porridges based on the state of the products are divided into two types: quick-cooking rice porridge and ready-to-eat rice porridge. The former can be served as rice porridge after rehydration in excess of hot or boiling water. This product has been developed and patented over the past years (Roberts, 1972; Luh and Liu, 1980; Robert et al., 1980; Juliano and Sakurai, 1985). However, the greatest disadvantage of this product is its relatively long preparation time compared to ready-to-eat rice porridge, which can be eaten after opening the package or heating.

Ready-to-eat rice porridge includes three kinds: plain rice porridge, savoury rice porridge and sweet rice porridge. As is well known, the preparation of plain rice porridge is rather simple, its essential ingredients are regular rice and water. However, study has yet to be conducted concerning the development of plain rice porridge into a commercial product suitable for local market.

Naivikul (1988) reported that glutinous rice tastes better than regular rice. It is often added into regular rice in small amounts to increase cooking and eating quality of the rice (Luh, 1991; Chen et al., 1995). Chang (1988) reported that rice 'congee' in Taiwan was all sweet, used glutinous rice as a base, and incorporated other ingredients. Other researchers also developed rice porridge using glutinous rice (Zhao, 1995; Yang, 1995). In Malaysia, there is a lot of broken rice from local milling and it is normally used for rice flour or other processed products such as vermicelli and noodles. The price of broken rice is very low, just a half of that of long-grain rice. Therefore, efforts should be made to determine the optimum composite rice based on the percentages of broken rice, glutinous rice and long-grain rice for producing plain rice porridge.

Plain rice porridge is less attractive to consumers because it is basically a bland food and low in nutritional value. Thus, it is often incorporated with ingredients containing dehydrated chicken, fish, beef, egg, legumes, vegetables and seasonings to compensate for the lack of nutrition and improve attractiveness to consumers. It can also be adapted to develop a wide range of products such as ready-to-eat savoury rice porridge by varying the ingredients used. Recently, the formulations of sweetened rice porridge were studied widely (Ding, 1995; Zhao, 1995; Yang, 1995). However, as reported by Jiang and Zhang (1995), the study on sweetened rice porridge was overwhelming and lesser has been conducted to study the formulation of savoury rice porridge. Therefore, it is also necessary to develop

ready-to-eat savoury rice porridges with fresh meats, fish, legumes, vegetables and seasonings to meet local people's preference for this kind of product.

A survey of food products available in the market showed that ready-to-eat nutritious food products for children are lacking. The recommended daily dietary allowances for 7-12 years old primary school children in Malaysia are 2150-2600 kcal, 35-54 g protein, 450-650 mg calcium and 10 mg iron (Siong et al., 1988). To meet this requirement, a child aged 7-12 years has to consume combinations of cereal, wheat, fish, egg and vegetables or fruit per meal. These combinations can be presented to a Malaysian child in the form of a rice-based dish. The rice-based dish, however, can be further categorised into boiled or steamed rice dish and rice porridge.

There is no standards available for the quality of rice porridge and the quality characteristics reported by various researchers were quite varied from each other (Ding, 1995; Zhao, 1995; Yang, 1995; Jiang and Zhang, 1995). In general, they evaluated the quality of rice porridge in terms of physicochemical, microbiological and sensory characteristics. Observations on the quality changes of ready-to-eat rice porridge on storage, however, have not been reported.

A study was therefore conducted with the following objectives:

1. To formulate canned plain rice porridge which is physically, organoleptically and economically acceptable
2. To formulate tasty, nutritious, and shelf-stable ready-to-eat canned savoury rice porridges for 7-12 years old children
3. To determine the quality changes of ready-to-eat canned savoury rice porridges during storage at different temperatures